

FURTHER MATHEMATICS/MATHEMATICS (ELECTIVE)

EXAMINATION SCHEME

There will be two papers, Papers 1 and 2, both of which must be taken.

PAPER 1: will consist of forty multiple-choice objective questions, covering the entire syllabus. Candidates will be required to answer all questions in $1\frac{1}{2}$ hours for 40 marks. The questions will be drawn from the sections of the syllabus as follows:

Pure Mathematics	-	30 questions
Statistics and probability	-	4 questions
Vectors and Mechanics	-	6 questions

PAPER 2: will consist of two sections, Sections A and B, to be answered in $2\frac{1}{2}$ hours for 100 marks.

Section A will consist of eight compulsory questions that are elementary in type for 48 marks. The questions shall be distributed as follows:

Pure Mathematics	-	4 questions
Statistics and Probability	-	2 questions
Vectors and Mechanics	-	2 questions

Section B will consist of seven questions of greater length and difficulty put into three parts: Parts I, II and III as follows:

Part I: Pure Mathematics	-	3 questions
Part II: Statistics and Probability	-	2 questions
Part III: Vectors and Mechanics	-	2 questions

Candidates will be required to answer four questions with at least one from each part for 52 marks.

SAMPLE QUESTIONS

PAPER 1 (OBJECTIVES)

- Find the equation of the line joining points (8, 1) and (-3, 4).
 - $3x - 11y - 35 = 0$
 - $3x - 11y + 35 = 0$
 - $3x + 11y - 35 = 0$
 - $3x + 11y + 35 = 0$
- The sum of the first and sixth terms of an Arithmetic progression (A.P.) is 21. If the first term is 3, find the eighth term.
 - 24
 - 27
 - 30
 - 33
- If α and β are the roots of the equation $2x^2 - 5x + m = 0$, where m is a constant, find $(\alpha^2 + \beta^2)$ in terms of m .
 - $\frac{25}{4} + 2m$
 - $\frac{25}{4} + m$
 - $\frac{25}{4} - 2m$
 - $\frac{25}{4} - m$
- Given that $y = \cos^2 x$, find $\frac{dy}{dx}$.
 - $-\sin^2 x$
 - $-\cos x \sin x$
 - $-2\cos x \sin x$
 - $-2\sin^2 x$

5. The position vectors of points P, Q and R are $\mathbf{p} = 4\mathbf{j}$, $\mathbf{q} = (4\mathbf{i} + 10\mathbf{j})$ and $\mathbf{r} = (k\mathbf{i} + 8\mathbf{j})$ respectively, where k is a constant. If $\angle PQR = 90^\circ$, find the value of k.
- A. 7
B. 1
C. -1
D. -7

PAPER 2
(ESSAY)

SECTION A

1. Given that * is a binary operation defined on R, the set of real numbers by $x*y = \frac{x^2}{x+y}$, where $x, y \in R$.
- (a) evaluate $(2 * 3) * 5$.
- (b) If $(x + 1) * (x + 2) = \frac{1}{3}$, find the value of x.
2. The marks scored by forty candidates in an examination are shown in the table.

Marks	1	2	3	4	5	6	7	8	9
Number of candidates	2	3	m	8	10	5	3	3	n

If the mean of the distribution is 4.725, find the values of m and n.

SECTION B


Part I
(Pure Mathematics)

3. Use the trapezium rule, with ordinates at $x = 1, 2, 3, 4$ and 5 , to calculate, correct to two decimal places, an approximate value for $\int_1^5 (2x + 8x^{-2}) dx$.

Part II
(Statistics and Probability)

4. The deviations from 10 of a given set of numbers are 2, 1, 0, -4, -5, -1, -2 and -7. Find the:
- (i) mean;
 - (ii) median;
 - (iii) standard deviation of the numbers.

Part III
Vectors and Mechanics



Coplanar forces 4N, 8N, 6N, 4N and 5N act at a point as shown in the diagram. If the 6N force act in the direction 090° , calculate the magnitude of the resultant force.